

論文

1. Prognostic Value of Angiographic Lesion Complexity in Patients With Acute Coronary Syndromes Undergoing Percutaneous Coronary Intervention (from the Acute Catheterization and Urgent Intervention Triage Strategy Trial). *Am J Cardiol*. 2014 (in press)
2. Impact of combined supine and prone myocardial perfusion imaging using an ultrafast cardiac gamma camera for detection of inferolateral coronary artery disease. *Int J Cardiol* 2014;174:313-7
3. Appearance of neointima according to stent type and restenotic phase: analysis by optical coherence tomography. *EuroIntervention* 2013;9:601-7
4. Impact of smoking on outcomes of patients with ST-segment elevation myocardial infarction (from the HORIZONS-AMI Trial). *Am J Cardiol* 2011 ;108:1387-94.
5. Prognostic significance of coronary thrombus in patients undergoing percutaneous coronary intervention for acute coronary syndromes: a subanalysis of the ACUITY (Acute Catheterization and Urgent Intervention Triage strategy) trial. *JACC Cardiovasc Interv* 2011;4:769-77.
6. Predictors of outcomes in medically treated patients with acute coronary syndromes after angiographic triage: an Acute Catheterization And Urgent Intervention Triage Strategy (ACUITY) substudy. *Circulation* 2010;121:853-62.
7. Pathological finding of sirolimus-eluting stent (SES) restenosis lesion with black hole appearance on intravascular ultrasound. *Circ J* 2009;73:1969-71.
8. Impact of impaired renal function and diabetes on long-term prognosis in patients undergoing primary angioplasty for acute coronary syndrome. *Intern Med* 2008;47:907-13.
9. Comparison of the incidence of acute coronary event after balloon angioplasty vs. stenting in native coronary arteries. *Intern Med* 2007;46:1501-6.